



Setzberg

Facts:

Town: **Spitz**
 Size in ha: **41,39**
 Altitude in m: **216 - 392**
 Aspect: **W - SE**
 Average Slope in %: **32**
 Max. Slope in %: **77**

Insolation in hours per year: **2101**
 Terraced vineyard: **Yes**
 Distance to Danube in m: **441**
 Labour required in h: **31.000**
 First documentary evidence: **1243**

Description

"The Setzberg must be divided into a large eastern half and a smaller western half. The probably most valuable dry grassland of the Wachau region lies between the two halves, which is also a designated natural monument.

In geological terms, the mountain unites several types of metamorphic rock: paragneiss, Spitzer granodiorite gneiss, marble, silicate marble, calc-silicate gneiss and scattered patches of loess on the east slope. In terms of the surface area, the variable paragneiss plays the most important role. It is therefore no surprise that the paragneisses occur in different forms on the Spitz and Laaben sides.

The location also represents a climatic divide separating the vineyards of Spitz from those of the Spitzer Graben. The wines produced on the two sides of the mountain differ accordingly."



Weingut Hirtzberger

Soil Profile Eastern Setzberg

The profile was taken from the highest vineyard on the Spitz side. The fine, sandy and stony soil, with humus accumulation in the upper 15 cm, lies directly above the compact parent rock consisting of paragneiss dominated by amphibolite. Fissures and joints are coated with carbonate precipitates, a weathering relict from the former loess cover.



Weingut Johann Donabaum

Soil Profile Western Setzberg

The profile from the Laaben side shows a sandy, stony topsoil on top of compact paragneiss interspersed with quartz veins. The rock is acidic, so both the fine earth and the cracks and joints are not influenced by loess and carbonates. The grapes grown here yield Rieslings rich in finesse.